

Survival of charmonia above T_c in anisotropic lattice QCD

Hideaki IIDA^{1,*}, Takumi DOI², Noriyoshi ISHII³, Hideo SUGANUMA⁴
and Kyosuke TSUMURA⁵

¹*YITP, Kyoto University, Sakyo, Kyoto 606-8502, Japan*

²*Dept. of Phys. & Astr., University of Kentucky, Lexington KY 40506, USA*

³*CCS, University of Tsukuba, Tsukuba, Ibaraki 305-8577, Japan*

⁴*Department of Physics, Kyoto University, Sakyo, Kyoto 606-8502, Japan*

⁵*Analysis Technology Center, Fujifilm Corporation, Kanagawa 250-0193, Japan*

We find a strong evidence for the survival of J/Ψ and η_c as spatially-localized $c\bar{c}$ (quasi-)bound states above the QCD critical temperature T_c , by investigating the boundary-condition dependence of their energies and spectral functions. In a finite-volume box, there arises a boundary-condition dependence for spatially spread states, while no such dependence appears for spatially compact states. In lattice QCD, we find almost *no* spatial boundary-condition dependence for the energy of the $c\bar{c}$ system in J/Ψ and η_c channels for $T \simeq (1.11 - 2.07)T_c$. We also investigate the spectral function of charmonia above T_c in lattice QCD using the maximum entropy method (MEM) in terms of the boundary-condition dependence. There is *no* spatial boundary-condition dependence for the low-lying peaks corresponding to J/Ψ and η_c around 3GeV at $1.62T_c$. These facts indicate the survival of J/Ψ and η_c as compact $c\bar{c}$ (quasi-)bound states for $T_c < T < 2T_c$.

§1. Introduction

J/Ψ suppression^{1),2)} was theoretically proposed as an important signal of quark-gluon plasma (QGP). The basic idea of J/Ψ suppression is that J/Ψ disappears above the QCD critical temperature T_c due to vanishing of the confinement potential and appearance of the Debye screening effect. In contrast, recently, some lattice QCD calculations indicate an opposite result that J/Ψ and η_c survive even above T_c .^{3),4),5),6),7),8)} Spectral functions of charmonia are extracted from temporal correlators at high temperature using the maximum entropy method (MEM) in Refs. 4)-8). Although there are some quantitative differences, the peaks corresponding to J/Ψ and η_c seem to survive even above T_c ($T_c < T < 2T_c$) in the $c\bar{c}$ spectral function.

However, one may ask a question on the “survival of J/Ψ and η_c ” observed in lattice QCD. Are the $c\bar{c}$ states above T_c observed in lattice QCD really compact (quasi-)bound states? Since colored states are allowed in the QGP phase, there is a possibility that the observed $c\bar{c}$ state in lattice QCD is just a $c\bar{c}$ scattering state,⁹⁾ which is spatially spread. Particularly in lattice QCD, even scattering states have discretized spectrum, due to the finite-volume effect. Therefore, for the fair judgment of the “survival of charmonia above T_c ”, it is necessary to clarify whether the $c\bar{c}$ systems are spatially compact (quasi-)bound states or scattering states.

To this end, we study the spatial *boundary-condition dependence* of the energy

*) E-mail: iida@yukawa.kyoto-u.ac.jp

and the spectral function for the $c\bar{c}$ systems (J/Ψ and η_c) above T_c in lattice QCD.⁷⁾ Actually, for $c\bar{c}$ scattering states, there occurs a significant energy difference between periodic and anti-periodic boundary conditions on the finite-volume lattice. In contrast, for spatially compact charmonia, there is almost no energy difference between these boundary conditions.⁷⁾ Using this fact, we investigate the $c\bar{c}$ system above T_c in terms of their spatial extent.

§2. Boundary-Condition Dependence and Compactness of States

We show briefly the method to distinguish spatially-localized states from scattering states on the finite-volume lattice.^{7),10)} For a compact $c\bar{c}$ (quasi-)bound state, the wave-function of the $c\bar{c}$ state is spatially localized, and therefore its energy is insensitive to the spatial boundary condition. In contrast, for a $c\bar{c}$ scattering state, the wave-function of the $c\bar{c}$ system is spatially spread, so that there occurs a significant boundary-condition dependence of the energy for the low-lying $c\bar{c}$ scattering state. Let us estimate the boundary-condition dependence.

- Under the *periodic boundary condition (PBC)*, the momentum of a quark or an anti-quark is discretized as $p_k = 2n_k\pi/L$ ($k = 1, 2, 3$, $n_k \in \mathbf{Z}$) on the finite lattice with the spatial volume L^3 , and the minimum momentum is $\vec{p}_{\min} = \vec{0}$.
- Under the *anti-periodic boundary condition (APBC)*, the momentum is discretized as $p_k = (2n_k + 1)\pi/L$ ($k = 1, 2, 3$, $n_k \in \mathbf{Z}$). In this case, the minimum momentum is $|\vec{p}_{\min}| = \sqrt{3}\pi/L$.

The energy difference of the low-lying $c\bar{c}$ scattering state is estimated as $\Delta E_{\text{scatt}} \equiv E_{\text{APBC}}^{\text{scatt}} - E_{\text{PBC}}^{\text{scatt}} \simeq 2\sqrt{m_c^2 + 3\pi^2/L^2} - 2m_c \simeq 350\text{MeV}$ for $L \simeq 1.55\text{fm}$ in Ref. 7), in the non-interacting case with the charm-quark mass m_c . In Ref. 7), we consider possible correction of the energy difference from a short-range potential of Yukawa type between a quark and an anti-quark, and find that the correction is negligible compared to the energy difference $\Delta E_{\text{scatt}} \simeq 350\text{MeV}$ estimated in the non-interacting case.

§3. Pole-Mass Measurements above T_c in Lattice QCD

First, we perform the standard pole-mass measurement of low-lying $c\bar{c}$ systems at finite temperature in anisotropic quenched lattice QCD⁷⁾ with the standard plaquette action at $\beta \equiv 2N_c/g^2 = 6.10$ and the renormalized anisotropy $a_s/a_t = 4.0$, i.e., $a_t = a_s/4 \simeq (8.12\text{GeV})^{-1} \simeq 0.024\text{fm}$ for the spatial and temporal lattice spacing. The adopted lattice sizes are $16^3 \times (14 - 26)$, which correspond to the spatial volume as $L^3 \simeq (1.55\text{fm})^3$ and the temperature as $(1.11 - 2.07)T_c$. We use 999 gauge configurations, picked up every 500 sweeps after the thermalization of 20,000 sweeps. For quarks, we use $O(a)$ -improved Wilson (clover) action on the anisotropic lattice.⁷⁾ We adopt the hopping parameter $\kappa = 0.112$, which reproduces the masses of charmonia at zero temperature. To enhance the ground-state overlap, we use a Gaussian spatially-extended operator with the extension radius $\rho = 0.2\text{fm}$ in the Coulomb gauge,⁷⁾ which is found to maximize the ground-state overlap. The energy of the low-lying $c\bar{c}$ state is extracted from the temporal correlator of the spatially-extended operators, where the total momentum of the system is projected to be zero.

Table I. The energy of the $c\bar{c}$ system in J/Ψ ($J^P = 1^-$) and η_c ($J^P = 0^-$) channels on PBC and APBC at $\beta = 6.10$ at each temperature. The superscripts J/Ψ and η_c denote quantities of J/Ψ and η_c , respectively. All the statistical errors are smaller than 0.01GeV. The energy difference $E_{\text{APBC}} - E_{\text{PBC}}$ observed in lattice QCD is also added. The observed energy difference is very small, compared to the estimated energy difference $\Delta E_{\text{scatt}} \simeq 350\text{MeV}$ in the case of the low-lying $c\bar{c}$ scattering states.

Temperature	$E_{\text{PBC}}^{J/\Psi}$	$E_{\text{APBC}}^{J/\Psi}$	$E_{\text{APBC}}^{J/\Psi} - E_{\text{PBC}}^{J/\Psi}$	$E_{\text{PBC}}^{\eta_c}$	$E_{\text{APBC}}^{\eta_c}$	$E_{\text{APBC}}^{\eta_c} - E_{\text{PBC}}^{\eta_c}$
$1.11T_c$	3.05GeV	3.09GeV	0.04GeV	3.03GeV	3.02GeV	-0.01GeV
$1.32T_c$	2.95GeV	2.98GeV	0.03GeV	2.99GeV	2.98GeV	-0.01GeV
$1.61T_c$	2.94GeV	2.98GeV	0.04GeV	3.00GeV	2.97GeV	-0.03GeV
$2.07T_c$	2.91GeV	2.93GeV	0.02GeV	3.01GeV	3.00GeV	-0.01GeV

Table I shows the boundary-condition dependence of the low-lying $c\bar{c}$ state energy in J/Ψ ($J^P = 1^-$) and η_c ($J^P = 0^-$) channels at finite temperatures. Both in J/Ψ and η_c channels, the energy difference between PBC and APBC is less than 40MeV, which is much smaller than the energy difference ΔE_{scatt} in the case of the low-lying $c\bar{c}$ scattering states, *i.e.*, $|E_{\text{APBC}} - E_{\text{PBC}}| \ll \Delta E_{\text{scatt}} \simeq 350\text{MeV}$, at all measured temperatures. These results indicate that the observed $c\bar{c}$ states are *spatially-localized (quasi-)bound states* as charmonia of J/Ψ and η_c for $1.11T_c < T < 2.07T_c$.

Here, we comment on a “constant contribution” to the meson correlator above T_c , provided by “wrap-around quark propagation”, which is pointed out in Ref. 11). The η_c channel is *free from* this extra contribution, but the J/Ψ channel suffers from it. Actually, in contrast to almost no temperature dependence of the η_c mass, there is a temperature dependence of the J/Ψ mass, which may be an artifact due to the constant contribution to the meson correlator.¹¹⁾ However, the difference of the J/Ψ mass between $1.11T_c$ and $2.07T_c$ is only about 140MeV, and this value is rather small compared to the estimated energy difference $\Delta E_{\text{scatt}} \simeq 350\text{MeV}$ for $c\bar{c}$ scattering states. Therefore, even including this extra effect, our main conclusion is unchanged also for J/Ψ above T_c . In fact, J/Ψ and η_c survive as spatially-localized (quasi-)bound states for $1.11T_c < T < 2.07T_c$.

§4. MEM Analyses for Spectral Functions above T_c in Lattice QCD

Next, we investigate the boundary-condition dependence of the spectral function $A(\omega)$ of the $c\bar{c}$ system above T_c using the maximum entropy method (MEM) in lattice QCD.⁷⁾ Using MEM, we extract the spectral function $A(\omega)$ from the temporal correlator $G(t)$ of the point source and the point sink, where the total momentum is projected to be zero. In this calculation, we use the Wilson quark action on a fine lattice at $\beta = 7.0$, *i.e.*, $a_t = a_s/4 \simeq (20.2\text{GeV})^{-1} \simeq 9.75 \times 10^{-3}\text{fm}$. The adopted lattice size is $20^3 \times 46$, which corresponds to the spatial volume $L^3 \simeq (0.78\text{fm})^3$ and the temperature $T \simeq 1.62T_c$.

Figure 2 shows the spectral functions in J/Ψ and η_c channels on PBC (dotted line) and APBC (solid line). There appear low-lying peaks around 3GeV, which correspond to the charmonia of J/Ψ and η_c . We find no spatial boundary-condition




Fig. 1. The spectral function $A(\omega)$ of the $c\bar{c}$ system in (a) J/Ψ channel and (b) η_c channel at $1.62T_c$ on PBC (dotted line) and APBC (solid line), normalized by the default function $m(\omega)$.⁷⁾ Almost no boundary-condition dependence is found for the low-lying peaks around 3GeV, which correspond to the charmonia of J/Ψ and η_c . In the η_c channel, the dotted and the solid lines coincide. For the figures of the spectral functions with errorbars, see Figs. 9 and 10 in Ref. 7).

dependence for the low-lying peaks, which indicate that the $c\bar{c}$ states corresponding to J/Ψ and η_c appear as spatially-localized (quasi-)bound states even above T_c .⁷⁾

§5. Summary and Conclusion

We have studied J/Ψ and η_c above T_c in anisotropic lattice QCD to clarify whether these states are spatially-localized (quasi-)bound states or $c\bar{c}$ scattering states. As a result, both in J/Ψ and η_c channels, we have found almost no spatial boundary-condition dependence of the energy of the low-lying $c\bar{c}$ system even on a finite-volume lattice for $(1.11 - 2.07)T_c$. Also in the MEM analysis, we find no spatial boundary-condition dependence of the low-lying peaks corresponding to J/Ψ and η_c in the spectral function at $T \simeq 1.62T_c$. These facts indicate that J/Ψ and η_c survive in QGP as spatially-localized (quasi-)bound states for $T_c < T < 2T_c$.

Acknowledgement

H. I. and H. S. thank the Yukawa Institute for Theoretical Physics, for fruitful discussions at “New Frontiers in QCD 2008”.

References

- 1) T. Hashimoto, O. Miyamura, K. Hirose and T. Kanki, Phys. Rev. Lett. **57** (1986), 2123.
- 2) T. Matsui and H. Satz, Phys. Lett. **B178** (1986), 416.
- 3) T. Umeda, K. Katayama, O. Miyamura and H. Matsufuru, Int. J. Mod. Phys. **A16** (2001), 2215; H. Matsufuru, O. Miyamura, H. Suganuma and T. Umeda, AIP Conf. Proc. **CP594** (2001), 258.
- 4) T. Umeda, K. Nomura and H. Matsufuru, hep-lat/0211003, Eur. Phys. J. **C39** (2005), 9.
- 5) M. Asakawa and T. Hatsuda, Phys. Rev. Lett. **92** (2004), 012001.
- 6) S. Datta, F. Karsch, P. Petreczky and I. Wetzorke, Phys. Rev. **D69** (2004), 094507.
- 7) H. Iida, T. Doi, N. Ishii, H. Suganuma and K. Tsumura, Phys. Rev. **D74** (2006), 074502; H. Iida, T. Doi, N. Ishii and H. Suganuma, PoS **LAT2005** (2005), 184.
- 8) G. Aarts *et al.* Phys. Rev. **D76** (2007), 094513.
- 9) Á. Mócsy and P. Petreczky, Phys. Rev. **D77** (2008), 014501.
- 10) N. Ishii, T. Doi, H. Iida, M. Oka, F. Okiharu and H. Suganuma, Phys. Rev. **D71** (2005),

- 034001; N. Ishii *et al.*, Phys. Rev. D**72** (2005), 074503.
11) T. Umeda, Phys. Rev. D**75** (2007), 094502.